Analysis of One-Year Cancer Survival Rates

NHS Brent CCG Profile



Transforming Cancer Services Team for London

TCST is part of the Healthy London Partnership, a collaboration between all London CCGs and NHS England London region to support the delivery of better health in London

Version:	3.7
Ref No:	P662
Date:	March 2016
Author:	PHAST Associates Paul Iggulden, Sarah Seager, Sue Atkinson
Commissioner:	Andy McMeeking, Team Manager – Transforming Cancer Services Team for London



www.phast.org.uk

Document Details								
Date of Issue	March 2016	Version	3.7					
The Public Health Action Support Team (PHAST) has undertaken this research. Transforming Cancer Services Team for London, commissioned the work.								
Commissioner Lead	Contact Details							
Name:	Andy McMeeking							
Role:	Team Manager – Transforming Cancer Services	Team for London						
Email:	andy.mcmeeking@nhs.net							





Table of Contents

Executive Summary	1
1. Introduction	5
2. The one-year cancer net survival index	7
2.1 Overview of section	7
2.2 Background to the one-year cancer net survival index	7
2.3 Overall all ages net survival measure	10
2.4 Site specific levels of survival: lung, colorectal and breast	12
2.4.1 Lung cancer	12
2.4.1 Colorectal Cancer	13
2.4.2 Breast Cancer	13
2.5 Age group levels of survival: all adults, 55-64ys and 75-99ys	15
2.6 One-year survival index for all cancers, all adults aged 15-99	16
2.7 One-year survival index for all cancers, adults aged 55-64	16
2.8 One-year survival index for all cancers, adults aged 75-99 years old	17
2.9 The one-year cancer net survival index: summary of section	18
3. The explanatory factors	20
3.1 An overview of the National Awareness and Early Diagnosis Initiative	20
3.2 Public awareness measures	21
3.2.1 Cancer Awareness Measures (CAM) survey	22
3.2.1.1 Barriers to seeking help	24





3.2.2 Healthy life expectancy at birth and 65 years
3.3 Early diagnosis measures
3.3.1 Screening measures; breast and bowel
3.3.1.1 Breast cancer screening
3.3.1.2 Bowel cancer screening
3.3.2 Use of the urgent referral pathway32
3.3.3 Emergency presentations
3.3.3.1 Proxy measure for emergency presentations for cancer
3.3.3.2 Rate of emergency admissions with cancer
3.3.4 Routes to diagnosis
3.3.5 Stage at diagnosis41
3.3.5.1 Prevalent cancer cases
3.3.5.2 Cancer Staging41
3.4 Diagnostics and treatment
3.4.1 6 week waits for key diagnostics43
3.4.2 Access to diagnostic procedures
3.4.3 Cancer waiting times
3.4.3.1 2 week wait, all cancer (patients waiting within 14 days)
3.4.3.2 31-day (diagnosis to treatment) wait for first treatment, all cancers
3.4.3.3 62-day (urgent GP referral) wait for first treatment, all cancer
3.4.4 Key provider performance





	3.5	Explanatory factors: summary of section	53
4.	Ne	ext steps	.56





Executive Summary

This report considers cancer survival and other measures related to cancer from a variety of statistics, for all cancers and for three main cancers - breast, lung and colorectal, and for different age groups. The results give an indication of how well Brent residents are being screened, diagnosed and treated for cancer in comparison with those in the rest of London and West Essex and in comparison with England averages. The findings indicate where the CCG may want to examine further or take action in their commissioning processes.

The index of one-year net survival from all cancers provides a convenient, single number that summarises the overall patterns of cancer survival, for a wide range of cancers with very disparate survival.

The methodology has been developed by the Cancer Survival Group (CSG) at London School of Hygiene and Tropical Medicine, this team being recognised as a centre of excellence within the UK and internationally, for the production of cancer survival statistics.

The all-cancers net survival index is adjusted for changes over time in the profile of cancer patients by age, sex and type of cancer. This is because survival varies widely with these three factors. The cancer net survival index is modelled to take into account likely mortality from non-cancer causes, using deprivation adjusted life tables.

The one-year survival index data is for the period 1997 to 2012, with sitespecific figures for the period 2008 to 2012. The explanatory measures presented follow these periods as closely as possible using the latest and most appropriate data available.

Taking this overall measure, the one-year net survival index for Brent has increased steadily to 70.2% of those with all newly diagnosed cancers





surviving one year or more in 2012 (latest data available from ONS), and is higher than the London average of 69.7% and the overall England figure of 69.3%¹. The 2012 measure shows Brent as having the one of the highest net survival rates amongst London and West Essex CCGs (ranking 11th; 1 highest, 33 lowest).

The following explanatory measures will have positively impacted on the level of one-year net survival from all cancers for Brent:

- The rate of emergency admissions with cancer per 100,000 population was lower for Brent than England as a whole for 2012 and 2013
- The proportion of colorectal cancer detected through screening is higher than the England average
- The latest staging data from the Cancer Outcomes and Services Dataset (COSD) shows similar percentages of cancers being diagnosed at each stage for Brent residents compared to those for London overall
- The proportion of the waiting list made up of people waiting more than six weeks for key diagnostic procedures tends to be lower in Brent than London overall
- In 2013/14, 96.1% of patients in NHS Brent CCG saw a specialist within two weeks. This is higher than the standard set and higher than the England average (95.3%)
- The percentage of non-small cell lung cancer (NSCLC) patients with cancer at stages 1a to 2b receiving surgery is considered a particular strong measure of provider performance and here North

¹ <u>http://www.ons.gov.uk/ons/rel/cancer-unit/a-cancer-survival-index-for-clinical-commissioning-groups/adults-diagnosed-1997-2012-and-followed-up-to-2013/index-of-cancer-survival-for-clinical-commissioning-groups.html#tab-Keypoints</u> last accessed 16 October 2015

West London Hospitals NHS Trust is higher than the London average (71.2% compared with 52%)

- The percentage of patients having CT prior to bronchoscopy is also considered a strong measure of provider performance and here North West London Hospitals NHST is higher than the London average (95.5% compared with 95%)
- The adjusted 18-month stoma rate is an area in which North West London Hospitals NHS Trust performs well.

The following explanatory measures will have negatively impacted on the level of one-year net survival from all cancers for Brent and suggest areas for improvement:

- For both males and females, and both at birth and at aged 65 years, healthy life expectancy in Brent is lower than that for England
- Breast cancer screening coverage and uptake in 2012 2014 were
 lower in Brent than the England average
- Brent residents in the 60-69 years target age group consistently have lower coverage and uptake for bowel cancer screening than for England as a whole
- The rate of urgent referrals per 100,000 population (referred using the two-week wait urgent referral pathway) was lower in Brent than England across the three years 2012, 2013 and 2014 and lower than a number of London CCGs
- The conversion rate was likewise lower for Brent at 9.1% in 2012 falling to 7.1% in 2014 against England figures of 10%. The conversion rate is the percentage of urgent referrals that were found to have cancer and provides an indicator of referral quality
- The detection rate tells us the percentage of all cancers that were identified through the two-week wait pathway. This was

approximately 47% in Brent, (in 2012 - 2014). This is below levels achieved by other London CCGs and the England average

- Approximately a quarter of Brent cancer patients presented as emergencies as measured by the NCIN proxy measure. The percentage has fluctuated around this level during 2008 - 2012 and is higher than that for England as a whole
- The proportion of breast cancer detected through screening is lower than the England average
- The proportion of colorectal cancer detected through the managed route is much lower than nationally and the proportion through emergency presentations much higher
- The prevalent cancer cases (% of practice population on practice cancer registers) for Brent residents in each of the three years 2012
 2014, was lower than the England value
- There are lower rates of access in Brent than in England for key diagnostic procedures and in the case of sigmoidoscopy in particular, these differences are more marked in 2013 than in 2012
- In 2013/14, 96.2% of cancer patients in NHS Brent CCG received their first treatment within 31 days of a decision to treat. Whilst slightly higher than the standard set this is lower than the England average (98.2%)
- In 2013/14, 82.2% of cancer patients in NHS Brent CCG received their first treatment within 62 days of an urgent GP referral. This is lower than both the standard set and the England average (85.8%)
- Adjusted 90-day post op mortality is a strong measure of colorectal cancer provider performance and here North West London Hospitals NHS Trust is considerably higher than the London average (13.7% compared with 4.6%).

1. Introduction

Cancer is the cause group responsible for the majority of avoidable deaths in England and Wales³. As such it is unsurprising that cancer indicators feature prominently in the national set of outcome indicators for Clinical Commissioning Groups (CCGs)⁴. Section 1 of the 2015/16 CCG Outcomes Indicator Set focuses on preventing people from dying prematurely and 8 of the 24 indicators relate to cancer. The primary focus of this report is two of these measures - one-year survival from all cancers and one-year survival from breast, lung & colorectal (bowel) cancers.

Clinical Commissioning Groups (CCGs) have been requested by NHE England (London) as part of the Key Lines of Enquiry (KLoE) to CCG narratives for 2015/16 operational plans to explain how they will track oneyear net cancer survival rates and to detail their plans for improving this key measure. This report has been produced to help the CCG Board to meet this request; it examines the one-year net survival index for adults in Brent alongside a range of explanatory factors, and identifies areas and actions for improvement.

The Transforming Cancer Services Team (TCST) for London has commissioned PHAST to report on, for each of the 33 London and West Essex CCGs, an in-depth analysis of current positions and trends in one-year cancer survival. A range of explanatory factors has been considered to identify areas for improvement, and where relevant, recommendations are made to enable CCGs to target interventions to improve cancer awareness and screening, diagnosis and treatment.

³ <u>http://www.ons.gov.uk/ons/rel/subnational-health4/avoidable-mortality-in-england-and-wales/2013/stb.html</u> last accessed 29 October 2015

⁴ <u>https://www.england.nhs.uk/wp-content/uploads/2012/12/ccg-ois-2015-glance.pdf</u>last accessed 29 October 2015

The report is in two parts:

- The one-year net survival index here we provide an overview of the index and report on current positions and trends across the 33 London and West Essex CCGs
- 2. The explanatory factors here we review how well Brent has performed across a wide range of measures (thirty-three). These measures cover the National Awareness and Early Diagnosis Initiative (NAEDI) pathway and are grouped into three themes: public awareness, early diagnosis and treatment.

The report uses routinely available data sources such as the National Cancer Intelligence Network (NCIN) GP practice profiles. London data is provided as a comparator alongside trend data for the CCG; where London figures were not available we have used published national data. The data periods used in the report were selected on the basis of a combination of timeliness and relevance to the periods covered by the net survival index (1997 – 2012 for the one-year survival index and 2008 – 2012 for the site specific indices).

2. The one-year cancer net survival index

2.1 **Overview of section**

CCGs have been requested by NHS England to explain how they will track one-year cancer survival and to detail their plans for improving this key measure. This part of the report provides an overview of the index and presents findings for London and West Essex CCGs. In this section we report on:

- Background to the one-year net survival index for all cancers combined
- Overall (all-ages) net survival
- Site specific levels of survival: lung, colorectal and breast
- Survival by age group: all adults (15-99 years), 55-64 years, 75-99 years
- Summary.

2.2 Background to the one-year cancer net survival index

The one-year cancer net survival index has been developed to provide a robust high-level summary measure. It is included in the CCG Outcomes Indicator Set⁵ and the All Party Parliamentary Committee on Cancer⁶ campaigned for it to be included in the Delivery Dashboard of the 2015/16 Clinical Commissioning Group (CCG) Assurance Framework⁷. Indeed, the NHS operating plan for CCGs (the Forward View into Action) includes the fundamental requirement for plans to improve early diagnosis for cancer and to track one-year cancer survival⁸.

The index of one-year survival from all cancers combined provides a convenient, single number that summarises the overall patterns of survival for

⁵<u>http://www.england.nhs.uk/resources/resources-for-ccgs/ccg-out-tool/ccg-ois/</u> last accessed 22 October 2015

⁶http://www.macmillan.org.uk/GetInvolved/APPG/APPG.aspx last accessed 22 October 2015 ⁷http://www.england.nhs.uk/commissioning/ccg-auth/ last accessed 16 October 2015 ⁸http://www.england.nhs.uk/wp-content/uploads/2014/12/forward-view-plning.pdf last accessed 16 October 2015

cancers newly diagnosed each calendar year, for a wide range of cancers with very disparate survival. Survival for most cancers is either stable or rising steadily from year to year¹. This trend is reflected in the values of the index in the tables produced by ONS. However, patterns of cancer occurrence by age, sex and type of cancer, can shift quite quickly over time, especially in small areas. The survival index is designed to reflect real progress (or otherwise) by providing a summary measure of cancer survival that adjusts for any such shifts. It is intended to change only if cancer survival itself actually changes and hence is a good measure. It is designed for long-term monitoring of progress in overall cancer survival.

Indicator values are published on the ONS website¹. The Cancer Survival Group (CSG) at LSHTM has developed the methodology for the indicators. ONS commissions cancer survival figures from the CSG, since this team is recognised as a centre of excellence within the UK for the production of cancer survival statistics.

To make figures from the past comparable with those for today and in the future, the all-cancers survival index is adjusted for changes over time in the profile of cancer patients by age, sex and type of cancer. This is because survival varies widely with all three factors.

Overall cancer survival can change simply because the profile of cancer patients changes, even if survival at each age, for each cancer and in each sex has not changed. This adjustment is made by using a weighted average of all the cancer survival estimates for each age, sex and cancer, using the proportions of cancer patients diagnosed in England and Wales during 1996– 99 in each age group, sex and type of cancer as the standard weights. All values of the cancer survival index, past and future, are adjusted using the same standard weights. This means that the cancer survival index is not affected by changes over time in the proportion of cancers of different lethality in either sex (for example, a reduction in lung cancer or an increase in breast cancer). Similarly, the index will be unaffected by a change in the age profile

8

of newly diagnosed cancer patients, or a shift in the proportion of a given type of cancer between men and women.

The cancer net survival index is modelled to take into account likely mortality from non-cancer causes. Adjustment of life tables for area deprivation takes an account of differences between areas in levels of relative deprivation.

The one-year survival index (%) for all cancers combined is available by calendar year of diagnosis, by CCG, Area Team and England. All adults (15 – 99 years) who were diagnosed with a first, primary, invasive malignancy were eligible for inclusion. Patients diagnosed with malignancy of the skin other than melanoma are excluded. Cancer of the prostate was also excluded from the index, because the widespread introduction of prostate-specific antigen (PSA) testing since the early 1990s has led to difficulty in the interpretation of survival trends (Pashayan et al., 2006).

Further details on the index methodology, published data and figures can be found on the <u>ONS website</u>.

2.3 **Overall all ages net survival measure**

The one-year all ages (15-99 years) net cancer survival measure for London CCGs has been reviewed and we have summarised the available data (1997 to 2012) in the following at a glance 'RAG rating' table.

The 'RAG rating' was developed by categorising the index values on the basis of comparison with the England value as being either:

- Red (the CCG value is low compared with England)
- Amber (the CCG value is broadly comparable with England)
- Green (the CCG value is better than that for England).

The detailed data on which the RAG rating table is based is available on the ONS website.

The table below gives a summary 'RAG rating' for the overall all adults oneyear net survival measure (15-99 years) alongside similar for 55-64 years and 75-99 years age groups.

CCG	15-99ys	55-64ys	75-99ys
Barking and Dagenham			
Barnet			
Bexley	ĺ		
Brent			
Bromley			
Camden			
Central London (Westminster)			
City and Hackney			
Croydon			
Ealing			
Enfield			
Greenwich			
Hammersmith and Fulham			
Haringey			
Harrow			
Havering			
Hillingdon			
Hounslow			
Islington			
Kingston			
Lambeth			
Lewisham			
Merton			
Newham			
Redbridge			
Richmond			
Southwark			
Sutton			
Tower Hamlets			
Waltham Forest			
Wandsworth			
West London			
West Essex			

Table 1: Summary of one-year net survival index by CCG

Legend

CCG consistently above England values CCG trend line crossing England trend line CCG consistently below England values



Underlying data source: London School of Hygiene and Tropical Medicine, published on ONS website.

The one-year net survival index for all cancers combined in Brent has increased over the period to 70.2% in 2012 (latest data available from ONS), and the published figures show this to be higher than the London average of 69.7% and the overall England figure of 69.3%. The 2012 measure shows Brent as having a net survival amongst the top third of London CCGs (ranking 11th in London; 1 highest, 33 lowest).

2.4 Site specific levels of survival: lung, colorectal and breast

The survival index will only change over time (or between areas) if the levels of survival for the particular cancer site or age group change over time (or between areas). In this section trends in the three main cancers are provided (lung, colorectal and breast).

These are chosen as between them they account for 39% of cancer mortality (based on 2012 <u>The 20 Most Common Causes of Cancer Death in 2012</u> <u>Number of Deaths per Year, All Ages, UK⁹</u>). Lung cancer is by far the most common cause of cancer death in the UK. More than one in five (22%) cancer deaths are from lung cancer. Bowel cancer is the second most common cause of cancer death (10%) and, breast cancer is the third most common cause of cancer deaths overall (7%).

2.4.1 Lung cancer

One-year net survival from lung cancer in Brent was consistently better than that for England as a whole during 2008-2012, where the differences in each year were statistically significant. The one-year lung cancer net survival for Brent increased, in line with the upward trend for England as a whole, to 35.7% in 2012 (latest data available from ONS) higher than the overall England figure of 33.4%. The 2012 measure shows Brent as having the 11th

⁹ <u>http://www.cancerresearchuk.org/health-professional/cancer-statistics/mortality/common-</u> <u>cancers-compared#heading-Zero</u> last accessed 16 October 2015

highest net survival rate amongst London CCGs (1 highest, 33 lowest).



Figure 1: One-year net survival index for lung cancer, aged 15-99 years

Source: London School of Hygiene and Tropical Medicine, published on ONS website.

2.4.2 Colorectal Cancer

One-year survival from colorectal cancer in Brent was, over the 2008 - 2012 period considered, also higher than that for England as a whole. The one-year colorectal cancer net survival for Brent increased, in line with the upward trend for England as a whole, to 79.2% in 2012 (latest data available from ONS) higher than the overall England figure of 77.3%. The 2012 measure shows Brent as having the 12th highest net survival rate amongst London CCGs (1 highest, 33 lowest).





2.4.3 Breast Cancer

One-year survival from breast cancer in Brent was, over the 2008 - 2012 period considered, higher than England as a whole. Net survival for Brent, in line with the trend for England as a whole, rose slightly over the period from 96.5% in 2008 to 97.1% in 2012 (latest data available from ONS) slightly higher than the overall England figure of 96.4%. The 2012 measure shows Brent as having the one of the highest net survival rates amongst London CCGs (ranking 5th in London; 1 highest, 33 lowest).

Source: London School of Hygiene and Tropical Medicine, published on ONS website.



Figure 3: One-year net survival index for breast cancer, aged 15-99 years

Source: London School of Hygiene and Tropical Medicine, published on ONS website.

2.5 Age group levels of survival: all adults, 55-64ys and 75-99ys

As stated previously, the survival index will only change over time (or between areas) if the levels of survival for a particular cancer or age group change over time (or between areas). In this section trends in all adults, in the 55-64 years age group and in the 75-99 years age groups are detailed.

The cancer one-year net survival publications include data on the specific age groups 55-64 years old and 75-99 years old alongside all adults (15-99 years). The 55-64 years age group is of particular interest as this is a key age group in terms of opportunities for maximising survival. More than a third of cancers are diagnosed in people aged 75 and over¹⁰ making this also a key age group for consideration.

¹⁰ <u>http://www.cancerresearchuk.org/health-professional/cancer-statistics/incidence#heading-</u> <u>Two</u>

2.6 **One-year survival index for all cancers, all adults aged 15-99**

The adult (15-99 years old) one-year net survival index for all cancers has risen steadily at England, regional (London) and local (Brent) levels. Nationally, one-year cancer net survival increased from 59.7% in 1997 to 69.3% in 2012, an increase of just under 10%. The improvement in Brent has been in line with that of England as a whole (and London overall), increasing from 59.7% in 1997 to 70.2% in 2012, an increase of 10.5%. *We can see from the following chart the percentage one-year net survival increasing steadily in Brent in line with that for England and London.*



Figure 4: One-year survival index for all cancers combined, aged 15-99 years

Source: London School of Hygiene and Tropical Medicine, published on ONS website.

2.7 One-year survival index for all cancers, adults aged 55-64

The one-year net survival index for all cancers has increased in the 55-64 years age group at England, regional (London) and local (Brent) levels. *The increase in the index for 55-64 year olds in Brent is very similar to that for England as a whole.* At England level, the index in the 55-64 years age group increased from 66.8% in 1997 to 76.7% in 2012 (up 9.9%). In Brent the index increased from 66.8% in 1997 to 75.9% in 2012 (up 9.1%).



Figure 5: One-year survival index for all cancers combined, aged 55-64 years

2.8 One-year survival index for all cancers, adults aged 75-99 years old

In older people (75 -99 years old) one-year net survival index has risen at England, regional (London) and local (Brent) levels. Nationally, one-year cancer net survival increased from 47.4% in 1997 to 57.2% in 2012, an increase of just under 10%. *The improvement in Brent has been greater than that for England as a whole (and London overall), increasing from 47.8% in 1997 to 61% in 2012, an increase of 13.2%.*

Source: London School of Hygiene and Tropical Medicine, published on ONS website.



Figure 6: One-year survival index for all cancers combined, aged 75-99

Source: London School of Hygiene and Tropical Medicine, published on ONS website.

The index for the 75-99 years age group has risen considerably faster in Brent than England and London as a whole between 2005 and 2012. There may be learning here for other CCGs?

The above charts suggest that the increase in the index for the 75-99 years age group in Brent has contributed to the increase seen in the index for all ages. This leads us to question whether similar improvements could be made in the 55-64 years age group.

2.9 The one-year cancer net survival index: summary of section

The index of one-year survival from all newly diagnosed cancers provides a convenient, single number that summarises the overall patterns of cancer survival in each calendar year, for a wide range of cancers with very disparate survival.

The all-cancers survival index is adjusted for changes over time in the profile of cancer patients by age, sex and type of cancer. This is because survival varies widely with all three factors. The cancer survival index is not affected by changes over time in the proportion of cancers of different lethality in either

sex (for example, a reduction in lung cancer or an increase in breast cancer). Similarly, the index will be unaffected by a change in the age profile of newly diagnosed cancer patients, or a shift in the proportion of a given type of cancer between men and women.

The one year net survival measure for Brent has increased steadily over the period to 70.2% in 2012 (latest data available from ONS), and is higher than the London average of 69.7% and the overall England figure of 69.3%. The 2012 measure shows Brent as having one of the highest net survival rates amongst London CCGs (ranking 11th; 1 highest, 33 lowest).

One year survival rates for the 'big' cancers have improved (2008-2012), are favourable compared with the overall England figure though are not consistent in terms of London ranking (lung 11th, colorectal 12th breast 5th across 33 London and West Essex CCGs.

The index for the 75-99 years age group in Brent increased considerably more than in England and London as a whole between 2005 and 2012.

3. The explanatory factors

In this section we review how well Brent has performed across a wide range of measures (thirty-three); these were chosen by the project team (PHAST and TCST) as they cover the 'NAEDI pathway' and are routinely available, making updating of the charts possible at a local level. This section presents:

- An overview of NAEDI and the NAEDI pathway
- Public awareness measures
- Early diagnosis measures
- Diagnostics and treatment measures

3.1 An overview of the National Awareness and Early Diagnosis Initiative

The National Awareness and Early Diagnosis Initiative (NAEDI) is a public sector/third sector partnership led by Cancer Research UK, the Department of Health, NHS England and Public Health England. NAEDI formally launched in 2008 and continues as a partnership between public and third sector organisations. Its role is to provide leadership and support to activities and research that promote earlier diagnosis of cancer.

When cancer is diagnosed at an early stage, treatment options, survival and chances of a full recovery are greater. Over 93% of bowel cancer patients diagnosed with the earliest stage of disease survive at least five years compared with less than 7% of those diagnosed with the most advanced stage disease¹¹. The same pattern is true for lung cancer, breast cancer, and for many cancers, common or rare. We know that inequalities exist, with some groups of patients more likely to be diagnosed with later stage disease.

11

http://www.cancerresearchuk.org/sites/default/files/health_professional_naedi_briefing_sheet. pdf last accessed 16th October 2015

The National Awareness and Early Diagnosis Initiative (NAEDI) was set out as a priority in the national strategy <u>Improving Outcomes: A Strategy for</u> <u>Cancer (January 2011)¹²</u>. The initiative promotes an evidence based pathway approach comprising of three key themes:

- Public awareness;
- Early diagnosis; and
- Diagnostics and treatment.

These three themes are used in the subsequent sections to structure our presentation of a wide range of measures that have been investigated to explain variation in the overall net survival measure.

3.2 **Public awareness measures**

When members of the public raise concerns and present possible symptoms to clinicians, the opportunities for early detection and treatment of less advanced cancers are increased. The National Awareness and Early Diagnosis Initiative (NAEDI) pathway starts with people knowing the signs of cancer and presenting to health services to have these investigated.

In this section we consider two important areas that will impact on early presentation of possible symptoms:

- Cancer awareness recall of a symptom of cancer (this provides a measure of knowledge of cancer symptoms) – and barriers to seeking help; and
- Healthy life expectancy (HLE).

¹² <u>https://www.gov.uk/government/publications/the-national-cancer-strategy</u> last accessed 16th October 2015

3.2.1 Cancer Awareness Measures (CAM) survey

Low cancer awareness may contribute to delayed presentation, delayed diagnosis and poor cancer survival. The Cancer Awareness Measure (CAM) is a validated survey that helps to assess people's knowledge of common cancer symptoms, and their intention to seek help if they notice a warning sign of cancer. In order to develop a baseline measure for the Cancer National Awareness and Early Diagnosis Initiative (NAEDI), the CAM was used in 23 of the 31 former PCT areas (London boroughs) between 2009 and 2011. Around 18,500 people were surveyed. We can see from the table below that survey results are known for two thirds of the 33 CCGs in London and West Essex.

In Brent, no CAM was conducted. Richmond achieved the highest percentage of respondents being able to recall a symptom of cancer (67%) and Enfield the lowest (30%).

	Unprompted recall of lump or swelling+	Year
England	68%	2008
Barking and Dagenham	31%	2009/10
Barnet	41%	2009/10
Bexley	-	-
Brent	-	-
Bromley	-	-
Camden	32%	2009/10
Westminster	49%	2009/10
City & Hackney	57%	2009/10
Croydon	60%	2009/10
Ealing	-	-
Enfield	30%	2009/10
Greenwich	55%	2012
Hammersmith and Fulham	-	2009/10
Haringey	35%	2009/10
Harrow	-	-
Havering	57%	2009/10
Hillingdon	*	2011
Hounslow	-	-
Islington	44%	2009/10
Kingston	64%	2009/10
Lambeth	57%	2009/10
Lewisham	-	-
Merton	54%	2009/10
Newham	48%	2009/10
Redbridge	50%	2009/10
Richmond	67%	2009/10
Southwark	-	-
Sutton	61%	2009/10
Tower Hamlets	42%	2009/10
Waltham Forest	48%	2009/10
Wandsworth	52%	2009/10
West Essex	41%	2009/10

Table 2: National and local Cancer Awareness Measures Survey results

+ in response to the question: "There are many warning signs and symptoms of cancer. Please name as many as you can think of" *Hillingdon, Bowel CAM only undertaken in 2011 Source: Materials collated by TCST London, September 2015.

The 2008 national survey of public awareness of cancer reported¹³ that "awareness of cancer warning signs was low when open-ended (recall) questions were used and higher with closed (recognition) questions; but on either measure, awareness was lower in those who were male, younger, and from lower socioeconomic status (SES) groups or ethnic minorities".

3.2.1.1 Barriers to seeking help

The report identified the most common barriers to seeking help were difficulty in making an appointment, worry about wasting the doctor's time and worry about what would be found. "Emotional barriers were more prominent in lower SES groups and practical barriers (e.g. too busy) more prominent in higher SES groups. Anticipated delay was lower in ethnic minority and lower SES groups".

Barriers to seeking help have been surveyed in national and local CAM surveys, with the following question posed:

"Sometimes people put off going to see the doctor, even when they have a symptom that they think might be serious. These are some of the reasons people give for delaying. Could you say if any of these might put you off going to the doctor?"

Findings from this area of the survey are presented in the following table:

¹³<u>http://www.cancerresearchuk.org/sites/default/files/bjc_awareness_in_britain_0.pdf</u> last accessed 16 October 2015

	Year	It would be difficult to make an appointment with my doctor	I would be worried about what the doctor might find	I would be too embarrassed	I would be worried about wasting the doctor's time	I would be too scared	I would be too busy to make time to go to the doctor	My doctor would be difficult to talk to	I have too many other things to worry about	I wouldn't feel confident talking about my symptom with the doctor	It would be difficult for me to arrange transport to the doctor's surgery	Number surveyed
England	2008	42%	37%	12%	38%	23%	32%	14%	24%	26%	4%	
All local CAM surveys*		28%	30%	17%	24%	22%	20%	13%	18%	11%	6%	49270
Barking & Dagenham	09/10	41%	30%	27%	22%	14%	19%	15%	18%	19%	7%	316
Barnet	09/10	21%	15%	16%	10%	24%	13%	5%	11%	5%	2%	737
Bexley	-	-	-	-	-	-	-	-	-	-	-	-
Brent	-	-	-	-	-	-	-	-	-	-	-	-
Bromley	-	-	-	-	-	-	-	-	-	-	-	-
Camden	09/10	15%	13%	11%	9%	9%	10%	5%	10%	7%	4%	555
Westminster	09/10	21%	11%	4%	10%	4%	22%	8%	17%	5%	4%	3053
City & Hackney	09/10	36%	33%	26%	33%	26%	36%	27%	29%	24%	12%	435
Croydon	09/10	30%	30%	13%	24%	22%	29%	14%	28%	14%	4%	835
Ealing	-	-	-	-	-	-	-	-	-	-	-	-
Enfield	09/10	20%	19%	10%	10%	16%	10%	10%	10%	6%	3%	644
Greenwich	2012	32%	30%	15%	12%	24%	24%	18%	20%	10%	6%	1230
Hammersmith & Fulham	09/10	-	-	-	-	-	-	-	I			-
Haringey	09/10	30%	34%	20%	20%	26%	22%	14%	21%	16%	11%	556
Harrow	-	-	-	-	-	-	-	-	-	-	-	-
Havering	09/10	33%	29%	18%	27%	14%	25%	21%	21%	17%	11%	513
Hillingdon	2011	22%	23%	9%	10%	13%	7%	10%	7%	11%	9%	338
Hounslow	-	-	-	-	-	-	-	-	-	-	-	-
Islington	09/10	23%	9%	2%	9%	11%	13%	6%	9%	4%	2%	449
Kingston	09/10	28%	27%	15%	25%	19%	29%	13%	26%	13%	6%	837
Lambeth	09/10	20%	29%	11%	15%	18%	17%	9%	14%	7%	5%	2224
Lewisham	-	-	-	-	-	-	-	-	-	-	-	-
Merton	09/10	33%	33%	15%	23%	23%	31%	17%	21%	17%	13%	834
Newham	09/10	38%	36%	31%	24%	21%	33%	21%	29%	20%	11%	359
Redbridge	09/10	38%	29%	24%	23%	18%	23%	27%	21%	21%	10%	542
Richmond	09/10	30%	25%	14%	27%	17%	33%	13%	21%	15%	6%	847
Southwark	-	-	-	-	-	-	-	-	-	-	-	-
Sutton	09/10	29%	36%	16%	24%	23%	30%	17%	29%	15%	4%	834
Tower Hamlets	09/10	43%	37%	26%	25%	20%	45%	29%	37%	24%	17%	368
Waltham Forest	09/10	22%	25%	18%	22%	17%	19%	14%	17%	16%	10%	463
Wandsworth	09/10	38%	34%	19%	28%	21%	44%	15%	39%	16%	9%	851
West Essex	09/10	15%	14%	18%	13%	20%	10%	8%	15%	10%	7%	630

Table 3: National and local Cancer Awareness Measures Survey results

Source: Materials collated by TCST London, September 2015.

National and local Cancer Awareness Measures Survey results

*Total refers to all local CAM surveys conducted in London.

The above table shows the variation of respondents being put off going to the doctor with symptoms because of perceived difficulties in making an appointment and being too embarrassed to go.

3.2.2 Healthy life expectancy at birth and 65 years

Healthy Life Expectancy (HLE) was selected because wellbeing in general and the absence of other medical conditions in particular, will tend to impact positively on the likelihood of people presenting with possible cancer symptoms; the hypothesis being that the presence of other conditions will mask cancer symptoms making it less likely for someone to recognise possible cancer symptoms and so delaying presentation of possible symptoms¹⁴. As such we might anticipate residents with better overall healthy life expectancy to present with possible cancer symptoms earlier and hence have a higher probability of surviving following cancer diagnosis.

Healthy life expectancy (HLE) is based on subjective self-assessed health and adds value to life expectancy by estimating the average lifetime spent in a favourable state of health. The following charts show healthy life expectancy (2010 - 2012) at birth and at 65 years old.

¹⁴<u>http://www.nature.com/bjc/journal/v113/n3/full/bjc2015164a.html</u> last accessed 16 October 2015



Figure 7: Healthy life expectancy at birth, 2010-12

Source: Office for National Statistics

Healthy life expectancy at age 65

At age 65 years old, men in England could expect to live another 9.2 years in "Good" health compared with 8.3 years for men in Brent. Women aged 65 years in England could expect to live 9.7 years in "good" health compared with 8.2 years for women in Brent.



Figure 8: Healthy life expectancy at age 65, 2010-12

For both males and females, and both at birth and at aged 65 years, healthy life expectancy in Brent is lower than that for England.

3.3 Early diagnosis measures

NAEDI highlights that various studies comparing cancer survival across countries show that England does worse than comparable countries¹⁰. With the exception of breast cancer, the gap in survival between England and other countries was not narrowing (when considering patients diagnosed up to 2007). Stage of the disease at diagnosis is a major factor in survival and, for some cancers there is evidence that more patients in England are diagnosed at a later stage compared with other countries¹⁰.

In this section we consider:

- 1. Screening measures; breast and bowel
- 2. Use of the urgent referral pathway
- 3. Emergency presentations
- 4. Routes to diagnosis
- 5. Stage at diagnosis.

Source: Office for National Statistics

3.3.1 Screening measures; breast and bowel

Figures compiled by Cancer Research UK show that approximately one third (34%) of cancer cases diagnosed in females aged 50-74 are breast cancers, many of which are diagnosed through screening¹⁵.

National screening programmes exist for specific age groups for breast cancer (women) and bowel cancer (men and women). In England, bowel screening has been shown to reduce the risk of dying from bowel cancer by a quarter in people who are screened¹⁶.

The national screening programmes are subject to on-going quality assurance to maximise effectiveness. Where there is variation in the impacts of these programmes this will likely reflect both how the service is delivered and patient/ population characteristics.

The measures included here detail:

- Coverage (percentage of target group screened in the recommended screening period e.g. 3 years for breast screening) and
- Uptake (percentage of target population screened within 6 months of invitation) for both breast and bowel cancer screening programmes.

3.3.1.1 Breast cancer screening

The one-year net breast cancer survival measure for Brent is higher than England as a whole and when compared to London (2012). This level of survival was amongst the top five London CCGs and was achieved even though breast cancer screening coverage and uptake in 2012 – 2014 were lower in Brent than the England average.

¹⁵<u>http://www.cancerresearchuk.org/health-professional/cancer-statistics/incidence/age#ref-6</u> last accessed 16 October 2015

¹⁶Hewitson P, Glasziou P, Irwig L, et al. <u>Screening for colorectal cancer using the faecal</u> <u>occult blood test (link is external)</u>. Hemoccult. Cochrane Database Syst Rev 2007(1): CD001216.





Source: NCIN, National GP Practice Profiles



Figure 10: Females aged 50-70 screened for breast cancer within 6 months of invitation

Source: NCIN, National GP Practice Profiles

3.3.1.2 Bowel cancer screening

The narrative for bowel cancer broadly echoes that for breast cancer. *Brent residents in the 60-69 years target age group consistently have lower coverage and uptake for bowel cancer screening than for England as a whole.*

Figure 11: Persons aged 60-69 screened for bowel cancer in last 30 months



Source: NCIN, National GP Practice Profiles





These high level measures might suggest further investigation in terms of effectiveness and service efficiency for breast and colorectal screening. Local quality assurance programme boards should be in place to support subsequent investigation as suggested by these high level measures.

3.3.2 Use of the urgent referral pathway

Cancer being diagnosed later is a major reason for poorer survival rates in the UK. The Improving Outcomes: A Strategy for Cancer⁸ estimates that, if patients were diagnosed at the same earlier stage as they are in other countries, up to 10,000 deaths could be avoided every year. The Strategy states that the challenge is clear: 'In order to improve early diagnosis, we need to encourage people to recognise the symptoms and signs of cancer and seek advice from their doctor as soon as possible. We also need doctors to recognise these symptoms and (if appropriate) refer people urgently for specialist care.' In a 2010 report, the National Audit Office identified that,

Source: NCIN, National GP Practice Profiles

amongst PCTs, there was almost a four-fold variation in the urgent cancer referral (two week wait) rate¹⁷.

Use of the urgent referral pathway is central to the NAEDI pathway and a key focus of service improvement initiatives. Here we present:

- The rate of urgent referrals per 100,000 population:(the 2 week wait urgent referral pathway)
- The proportion of patients referred who are subsequently diagnosed with cancer (conversion rate); and
- The proportion of cancer cases in the practice that were referred through the two-week wait route (detection rate).

To better understand variations in use of the urgent referral pathway and to achieve earlier diagnosis, it is important that all three of these measures are considered together along with measures of the use of diagnostic tests.

The rate of urgent referrals per 100,000 population (referred using the twoweek wait urgent referral pathway) was lower in Brent than England across the three years 2012, 2013 and 2014 and lower than a number of London Boroughs.

We see this in the following charts for overall referrals and for each of the three main cancer site-specific referrals, noting that lung cancer referrals have increased and in 2014 the rate is closer to the England rate.

¹⁷<u>http://www.nao.org.uk/wp-content/uploads/2010/11/1011568.pdf</u> last accessed 16 October 2015



Figure 13: Overall two-week-wait referrals

Source: NCIN, National GP Practice Profiles





Source: NCIN, National GP Practice Profiles



Figure 15: Two-week-wait referrals with suspected lower GI cancer

Source: NCIN, National GP Practice Profiles



Figure 16: Two-week-wait referrals with suspected lung cancer

Source: NCIN, National GP Practice Profiles

The conversion rate was likewise lower for Brent at 9.1% in 2012 falling to 7.1% in 2014 against England figures of 10%. The conversion rate is the percentage of the referrals that were found to have cancer and provides an indicator of the referral quality.





The detection rate tells us the percentage of all cancers that were identified through the two-week wait pathway. This was approximately 47% in Brent, (in 2012 - 2014). This is below levels achieved by other London Boroughs and the England average.





Source: NCIN, National GP Practice Profiles

Source: NCIN, National GP Practice Profiles

The two-week wait referrals measures suggest this is an area in need of improvement for Brent. The relatively low performance in this area suggests scope for GP training and support to improve the 2ww referral, conversion and detection rates

3.3.3 Emergency presentations

Emergency presentations suggest late presentation and diagnosis and hence reduced chance of survival. High levels of emergency presentation indicate the need for improved awareness raising and support for the public to recognise and act on possible symptoms, and/or training for professionals (GPs) in symptom recognition and earlier referral. There is clearly a link between this and the previous section on two-week wait referrals and this theme is explored in this section and the following on 'routes to diagnosis'.

This section includes:

- The National Cancer Intelligence Network (NCIN)¹⁸ proxy measure for emergency presentations for cancer; and
- the overall rate of emergency admissions.

3.3.3.1 Proxy measure for emergency presentations for cancer

The percentage of emergency presentations for cancer for Brent residents is broadly comparable with that for England as a whole. The following chart shows approximately a quarter of Brent cancer patients presenting as emergencies. The percentage has fluctuated around this level during 2008 -2012 and is higher than that for England as a whole.

¹⁸ National Cancer Intelligence Network, http://www.ncin.org.uk/cancer_information_tools/cct



Figure 19: Proxy measure for emergency presentations for cancer

Source: NCIN, National GP Practice Profiles

3.3.3.2 Rate of emergency admissions with cancer

The rate of emergency admissions with cancer per 100,000 population was lower for Brent than England as a whole for 2012 and 2013. In keeping with the rate for England overall, there was a large drop in the rate in Brent from 2012 to 2013. For both of these years, the rate in Brent was lower than that for England (figure 20).



Figure 20: Rate of emergency admissions with cancer



3.3.4 Routes to diagnosis

The routes to diagnosis produced by the National Cancer Intelligence Network provide a valuable summary picture of the route patients take to diagnosis. The following figures for female breast, colorectal and lung cancer develop further some of the key findings from the previous three sections, namely:

- Lower proportions detected through screening in Brent than England
- Lower proportions detected through managed care routes (such as the use of the urgent referral pathway).

The data on emergency presentations gives valuable context and additional detail to the overall emergencies presentations measures in 3.3.3 above. In particular we can see from these figures (below) that:

- The proportion of breast cancer detected through screening is lower than the England average
- Whilst the proportion of colorectal cancer detected through screening is higher than the England average, the percentage detected through the

managed route is much lower than nationally and the proportion through emergency presentations much higher.

There is a need to better understand the 'other' routes to diagnosis category; it has been suggested this might be the result of large numbers of patients being screened under private health care schemes.

The CCG is encouraged to explore means of reducing the proportion of breast and lung cancer cases detected as emergency presentations through improvements in screening and use of the urgent referral pathway.

Figure 21: Directly age-standardised rate per 100,000 population by route to diagnosis – breast cancer

Directly age-standardised rate per 100, 000 population by route Bordered cells indicate 3 SD outliers on respective funnel plots							Percentage by Route			
Female Breast Cancer	Screen detected	Managed	Emergency presentation	Other	Number of cases	Screen detected		Managed	Emergency presentation	Other
England	38.4	71.2	3.9	6.0	101 120	28	%	62%	5%	5%
Confidence interval	38.1 38.7	70.8 71.6	3.8 4.0	5.9 6.1	151,120	28%	29%	62% 62%	5% 5%	5% 5%
NHS Brent CCG	25.6	54.2	3.7	8.9	640	24	%	62%	5%	10%
Confidence interval	21.5 29.6	48.9 59.6	2.4 5.0	6.7 11.1	0.40	21%	27%	58% 65%	3% 7%	8% 12%

Source: NCIN

Figure 22: Directly age-standardised rate per 100,000 population by route to diagnosis – colorectal cancer

Directly age-standardised rate per 100, 000 population by route Bordered cells indicate 3 SD outliers on respective funnel plots							Percentage by Route			
Colorectal	Screen detected	Managed	Emergency presentation	Other	Number of cases	Screen detected	Managed	Emergency presentation	Other	
England	2.5	29.3	10.2	1.9	156 057	5%	66%	25%	4%	
Confidence interval	2.5 2.6	29.1 29.4	10.1 10.3	1.9 1.9	130,037	5% 5	% 66% 66%	25% 25%	4% 4%	
NHS Brent CCG	2.8	19.2	10.9	1.9	490	7%	55%	33%	5%	
Confidence interval	1.8 3.7	16.9 21.6	9.2 12.6	1.1 2.6	400	5% 10	1% 51% 59%	29% 37%	4% 8%	

Source: NCIN

Figure 23: Directly age-standardised rate per 100,000 population by route to diagnosis – lung cancer



Source: NCIN

3.3.5 Stage at diagnosis

The earlier cancer is detected the better the chance to improve survival. In this section we consider two measures:

- The proportion of the practice list on the practice cancer register (prevalent cancer cases); and
- Stage at diagnosis.

The first of these can be considered to be indicative of practice detection of cancers. Where a CCG has low percentages of residents on practice cancer registers it is anticipated that detection will be later and diagnosis will likely be at a later stage.

3.3.5.1 Prevalent cancer cases

The prevalent cancer cases (% of practice population on practice cancer registers) for Brent residents in each of the three years 2012 - 2014, was lower than the England value.



Figure 24: Prevalent cancer cases

Source: NCIN, National GP Practice Profiles

3.3.5.2 Cancer Staging

High-quality, comparable staging data on more than 80% of all cases of the most common cancers is now available for the whole of the country, making

England's National Cancer Registration Service one of the most advanced anywhere in the world.

The latest staging data from the Cancer Outcomes and Services Dataset (COSD)¹⁹ shows similar percentages of cancers being diagnosed at each stage for Brent residents compared to those for London overall.

These findings though should be treated with caution given the high proportions of cancers that were unknown (32% of staging data not submitted by the hospital trust) or unstageable (5%)* at diagnosis.



Figure 25: Proportions of patients at each stage of diagnosis, 2013

Source: COSD

3.4 **Diagnostics and treatment**

A range of patient and service factors will affect stage at diagnosis. We have considered above patient knowledge and GP referrals (two-week wait). Once

¹⁹ <u>http://www.ncin.org.uk/collecting_and_using_data/data_collection/cosd</u> last accessed 16 October 2015

^{*} Unstageable should be assigned only after all efforts to identify the extent of the disease have been exhausted or the site or histology does not meet criteria for staging. Some cancers such as blood cancer are not stageable.

referred, the time patients wait for key diagnostics will have a bearing on the stage at diagnosis.

In this section we review:

- 6 week waits for key diagnostics;
- Access to diagnostics;
- Cancer waiting times; and
- Key provider treatment performance measures.

3.4.1 6 week waits for key diagnostics

Colonoscopy, flexible sigmoidoscopy and gastroscopy are key procedures used to identify bowel and gastrointestinal cancers (diagnostic) or treat conditions (therapeutic). We see from the following charts that although the values are variable, the proportion of the waiting list made up of people waiting more than six weeks for key diagnostic procedures tends to be lower in Brent that London overall.



Figure 26: Patients waiting over 6 weeks for a colonoscopy

Source: http://www.england.nhs.uk/statistics/diagnostics-waiting-times-and-activity/



Figure 27: Patients waiting over 6 weeks for a flexi sigmoidoscopy

Source: http://www.england.nhs.uk/statistics/diagnostics-waiting-times-and-activity/

Figure 28: Patients waiting over 6 weeks for a gastroscopy



Source: http://www.england.nhs.uk/statistics/diagnostics-waiting-times-and-activity/

3.4.2 Access to diagnostic procedures

Here we review rates of key diagnostic procedures. In all three procedures there are lower rates of access in Brent than for England and in the case of

sigmoidoscopy in particular these differences are more marked in 2013 than in 2012.



Figure 29: In-patient or day-case colonoscopy procedures

Source: NCIN, National GP Practice Profiles



Figure 30: In-patient or day-case sigmoidoscopy procedures

Source: NCIN, National GP Practice Profiles



Figure 31: In-patient or day-case upper GI endoscopy procedures

Source: NCIN, National GP Practice Profiles

These measures suggest urgent action is required by Brent commissioners to improve patient waiting times and access to these procedures.

3.4.3 Cancer waiting times

The time taken for a patient to be diagnosed and for treatment to begin will impact on chances of survival. In this section we review the following measures:

- 2 week wait:
- 31 day wait; and
- 62 day wait.

3.4.3.1 2 week wait, all cancer (patients waiting within 14 days)

This measure is key in terms of early diagnosis. The standard set for England is 93%. In 2013/14, 96.1% of patients in NHS Brent CCG saw a specialist within two weeks. This is higher than the standard set and higher than the England average (95.3%).





Source: Cancer Waiting Times, NHS England

3.4.3.2 31-day (diagnosis to treatment) wait for first treatment, all cancers

This measure records patients receiving their first treatment within 31 days of the decision to treat. The standard set in England is 96%. We can see from the following chart that in 2013/14, 96.2% of cancer patients in NHS Brent CCG received their first treatment within 31 days of a decision to treat. Whilst slightly higher than the standard set this is lower than the England average (98.2%).



Figure 33: 31 day wait for first treatment

Source: Cancer Waiting Times, NHS England

3.4.3.3 62-day (urgent GP referral) wait for first treatment, all cancer

This measures the percentage of patients receiving their first treatment within 62 days of an urgent GP referral. The operational standard in England is 85%. In 2013/14, 82.2% of cancer patients in NHS Brent CCG received their first treatment within 62 days of an urgent GP referral. This is lower than both the standard set and the England average (85.8%).



Figure 34: 62 day urgent GP referral for first treatment

Source: Cancer Waiting Times, NHS England

3.4.4 Key provider performance

In this section we provide results from the 2014 National Lung and Colorectal cancer audit ²⁰. Whilst the focus of the audit is provider performance, rather than specifically the treatment for the CCGs residents, we can gain insights for Brent residents by focussing on the main provider for Brent, namely North West London Hospitals NHS Trust.

The following table shows performance at North West London Hospitals NHS Trust to be fairly good (green ratings in 5 of the 6 measures). The percentage of non-small cell lung cancer (NSCLC) patients with cancer at stages 1a to 2b receiving surgery is considered a particular strong measure of provider performance and here North West London Hospitals NHS Trust is higher than the London average (71.2% compared with 52%). The percentage of patients having CT prior to bronchoscopy is also considered a strong measure of

²⁰ National Bowel Cancer Audit: <u>http://www.hscic.gov.uk/bowel</u> National Lung Cancer Audit: http://www.hscic.gov.uk/lung

provider performance and here North West London Hospitals NHS Trust is higher than the London average (95.5% compared with 95%).

Table 4: Resu	lts by provider	from the 20	14 National	Lung Cancer	Audit,
London Cance	er and London	Cancer Alli	ance		

	% of discussed at patients MDT	% having CT prior to bronchoscopy	% of patients with a histological /cytological diagnosis	% CNS present at diagnosis	% of NSCLC 1a to 2b Receiving surgery 2011-13	% receiving Active treatment
Barking, Havering and Redbridge	88.6%	74.2%	80.4%	52.9	35.2%	50.4
Barnet and Chase Farm	95.6%	77.0%	91.3%	92.3	63.2%	72.1
Bart's Health/Newham	97.2%	82.9%	88.9%	66.7	41.7%	59.7
Bart's Health/RLH	96.7%	95.3%	90.9%	59.5	62.2%	58.7
Bart's Health/Whipps Cross	91.8%	90.5%	86.1%	77.0	45.2%	68.9
Chelsea and Westminster	94.4%	90.9%	91.7%	58.3	59.5%	61.1
Croydon Healthcare NHS Trust	97.0%	92.5%	85.1%	96.7	39.7%	61.2
Ealing Hospital NHS Trust	98.5%	82.4%	55.4%	66.2	31.0%	46.2
Epsom and St Helier Uni. NHS	76.2%	92.5%	73.5%	40.9	32.9%	59.7
Guys and St Thomas' NHS FT	100.0%	94.8%	100.0%	55.0	56.5%	87.4
Imperial College Healthcare NHS	100.0%	95.9%	86.7%	51.8	57.5%	73.1
Kings College Hospital NHS FT	99.3%	88.1%	84.7%	86.1	56.7%	61.3
Kingston Hospital NHS Trust	95.9%	90.9%	76.2%	59.0	60.0%	60.7
Lewisham	97.7%	79.1%	78.3%	67.1	44.2%	58.9
North Middlesex	97.8%	100.0%	80.0%	97.8	45.9%	64.4
North West London	96.8%	95.5%	86.2%	63.8	71.2%	67.0
South London Healthcare	95.9%	91.1%	81.8%	76.5	36.8%	62.4
St Georges Healthcare NHS	84.6%	81.8%	83.8%	35.3	53.7%	56.6
The Hillingdon NHS Foundation	93.5%	97.9%	58.1%	79.0	51.4%	52.4
The Homerton	100.0%	94.7%	83.7%	77.2	44.3%	59.8
The Princess Alexandra	99.4%	93.7%	81.6%	88.6	63.0%	64.6
The Royal Brompton and	21.7%	95.8%	100.0%	56.5	86.4%	87.0
The Royal Free	100.0%	96.2%	90.3%	97.1	79.6%	63.1
The Whittington	97.8%	96.3%	76.1%	37.0	51.0%	65.2
UCLH	95.6%	94.4%	88.5%	55.0	59.8%	85.5
West Middlesex University NHS	98.9%	97.3%	78.7%	1.1%	54.9%	45.7
London TOTAL	95.0%	95.0%	75.0%	80.0	52.0%	60.0

Source: National Lung Cancer Audit: <u>http://www.hscic.gov.uk/lung</u>

The following tables show colorectal cancer performance at North West London Hospitals NHS Trust (table 6) to be above average (green ratings in 4 of the 6 measures). The adjusted 90-day post op mortality is strong measure of colorectal cancer provider performance and here North West London Hospitals NHS Trust is considerably higher than the London average (13.7% compared with 4.6%). The adjusted 18-month stoma rate is an area in which North West London Hospitals NHS Trust performs well with a rate of 29% compared with a London audit average of 51%.

	Adjusted 90 day post op Mortality	% LOS > 5 Days	% of operations attempted laparoscopically	% who underwent major resection	Adjusted 2 year mortality	Adjusted 18 month stoma rate
Barnet and Chase Farm	3.7%	68.6%	67.8%	64.0%	18.6%	62.0%
Bart's Health	4.6%	75.2%	85.0%	65.5%	33.4%	46.0%
Barking, Havering & Redbridge NHST	6.8%	73.5%	52.9%	61.5%	19.1%	54.0%
North Middlesex	4.6%	90.9%	84.1%	68.8%	36.1%	0.0%
The Homerton	5.2%	84.4%	52.2%	64.8%	23.4%	28.0%
The Princess Alexandra	8.3%	42.9%	74.1%	74.2%	32.1%	44.0%
The Whittington	4.4%	78.7%	71.4%	60.0%	20.2%	45.0%
The Royal Free	8.8%	74.1%	53.2%	67.4%	Not reported	39.0%
UCLH	10.1	80.6%	58.0%	66.3%	25.2%	62.0%
London Cancer	5.6%	75.6%	67.8%	64.7%	24.4%	49.0%
Audit Average	4.6%	69.0%	61.0%	66.0%	24.0%	51.0%

Table 5: Results by provider from the 2014 National Colorectal CancerAudit, London Cancer

Source: National Bowel Cancer Audit: <u>http://www.hscic.gov.uk/bowel</u>

	Adjusted 90 day post op Mortality	% LOS > 5 Days	% of operations attempted laparoscopically	Adjusted 90 day unplanned readmission rate %	Adjsuted 2 year mortality	Adjusted 18 month stoma rate using HES
Chelsea and Westminster	5.6%	87.0%	47.9%	26.7%	19.4%	56.0%
Croydon Healthcare NHS Trust	9.9%	Data not submitted	41.3%	8.1%	21.0%	38.0%
Ealing Hospital NHS Trust	1.9%	90.2%	73.8%	31.0%	22.1%	53.0%
Epsom and St Helier University NHS Trust	5.2%	82.1%	82.1%	20.1%	24.0%	33.0%
Guys and St Thomas' NHS FT	3.8%	Data not submitted	100.0%	21.3%	17.5%	74.0%
Imperial College Healthcare NHS Trust	0.8%	95.0%	83.3%	25.4%	19.5%	51.0%
Kings College Hospital NHS FT	4.3%	80.9%	77.1%	14.3%	18.8%	61.0%
Kingston Hospital NHS Trust	7.8%	72.3%	36.6%	25.7%	22.1%	32.0%
Lewisham	0.0%	69.4%	18.2%	12.0%	30.0%	55.0%
North West London	13.7%	100.0%	83.3%	14.6%	18.3%	29.0%
St Georges Healthcare NHS Trust	4.6%	61.5%	65.1%	22.1%	n/a	26.0%
South London Healthcare	4.3%	71.9%	61.3%	19.8%	26.0%	31.0%
The Hillingdon NHS FT	2.0%	78.0%	58.3%	31.3%	31.7%	61.0%
The Royal Marsden	0.0%	85.7%	52.9%	10.2%	7.9%	25.0%
West Middlesex University NHS Trust	5.2%	67.3%	51.0%	22.7%	21.6%	46.0%
LCA Total	4.3%	78.8%	52.7%	20.6%	22.0%	54.0%
Audit Average	4.6%	69.0%	61.0%	19.8%	24.0%	51.0%

Table 6: Results by provider from the 2014 National Colorectal CancerAudit, London Cancer Alliance

Source: National Bowel Cancer Audit: <u>http://www.hscic.gov.uk/bowel</u>

3.5 **Explanatory factors: summary of section**

Here we provide a recap of the key points raised in part 2 of this report.

- For both males and females, at birth and at aged 65 years, healthy life expectancy in Brent is lower than that for England
- Breast cancer screening coverage and uptake in 2012 2014 were lower in Brent than the England average
- Brent residents in the 60-69 years target age group consistently have lower coverage and uptake for bowel cancer screening than for England as a whole
- The rate of urgent referrals per 100,000 population (referred using the two-week wait urgent referral pathway) was lower in Brent than England across the three years 2012, 2013 and 2014 and lower than a number of London CCGs
- The conversion rate was likewise lower for Brent at 9.1% in 2012 falling to 7.1% in 2014 against England figures of 10%. The conversation rate is the percentage of the referrals that were found to have cancer and provides an indicator of the referral quality
- The detection rate tells us the percentage of all cancers that were identified through the two-week wait pathway. This was approximately 47% in Brent, (in 2012 - 2014). This is below levels achieved by other London CCGs and the England average
- Approximately a quarter of Brent cancer patients presented as emergencies as measured by the NCIN proxy measure. The percentage has fluctuated around this level during 2008 - 2012 and is higher than that for England as a whole
- The rate of emergency admissions with cancer per 100,000 population was lower for Brent than England as a whole for 2012 and 2013
- The proportion of breast cancer detected through screening is lower than the England average

- The proportion of colorectal cancer detected through screening is higher than the England average
- The proportion of colorectal cancer detected through the managed route is much lower than nationally and the proportion detected through emergency presentations much higher
- The prevalent cancer cases (% of practice population on practice cancer register) for Brent residents in each of the three years 2012 -2014, was lower than the England value
- The latest staging data from the Cancer Outcomes and Services Dataset (COSD) shows similar percentages of cancers being diagnosed at each stage for Brent residents compared to those for London overall
- The proportion of the waiting list made up of people waiting more than six weeks for key diagnostic procedures tends to be lower in Brent that London overall
- There are lower rates of access in Brent than in England for key diagnostic procedures and in the case of sigmoidoscopy in particular these differences are more marked in 2013 than in 2012
- In 2013/14, 96.1% of patients in NHS Brent CCG saw a specialist within two weeks. This is higher than the standard set and higher than the England average (95.3%)
- In 2013/14, 96.2% of cancer patients in NHS Brent CCG received their first treatment within 31 days of a decision to treat. Whilst slightly higher than the standard set this is lower than the England average (98.2%)
- In 2013/14, 82.2% of cancer patients in NHS Brent CCG received their first treatment within 62 days of an urgent GP referral. This is lower than both the standard set and the England average (85.8%)
- The percentage of non-small cell lung cancer (NSCLC) patients with cancer at stages 1a to 2b receiving surgery is considered a particular strong measure of provider performance and here North West London

Hospitals NHS Trust is higher than the London average (71.2% compared with 52%)

- The percentage of patients having CT prior to bronchoscopy is also considered a strong measure of provider performance and here North West London Hospitals NHS Trust is higher than the London average (95.5% compared with 95%)
- The adjusted 90-day post op mortality is a strong measure of colorectal cancer provider performance and here North West London Hospitals NHS Trust is considerably higher than the London average (13.7% compared with 4.6%)
- The adjusted 18-month stoma rate is an area in which North West London Hospitals NHS Trust performs well.

4. Next steps

In this review we have examined over two dozen measures related to survival, screening, diagnosis and management of cancer patients, in particular the three main cancers - breast, colorectal and lung.

The key findings and suggested actions are detailed in the report and repeated in summary sections (at the end of Parts 1 and 2) and additionally in the Executive Summary.

It is hoped that CCGs will be engaged through this report and encouraged to consider what can be done to improve further survival from cancer. More detailed data may be available to enable supplementary analyses and TCST for London will assist with this as resources permit.

Finally, the TCST for London welcome your feedback on this report.

www.phast.org.uk

