

## Remember...

Some inhalers must be used with a spacer.  
Check with your GP, asthma nurse or pharmacist

Always keep your rescue inhaler and your spacer with you. You might need them if your asthma gets worse

Make sure you have an asthma review within 48 hours after an attack

## Extra Advice from my Asthma Professional:

## Additional Resources:

[Asthma and Lung UK Asthma Toolkit](#)

[Check you're using your inhaler properly:](#)



## My Asthma Triggers

### List the things that make your asthma worse:

Pollen	Vaping
Dust	Environmental pollution
Animal fur	Other fumes/ sprays
Weather	Respiratory infections (cold/flu)
Exercise	Medicines
Mould/damp	Stress/emotions
Fumes	Food *
Tobacco smoke	
House dust mite	

\* Always refer to your Allergy Plan as well  
**Any Other Triggers:**

## Contact Details

GP:

Asthma Specialist/Team:

This plan was approved by  
London Asthma Leadership and Implementation Group (LALIG)  
Approval Date: May 2024  
To Be Revised: May 2026

## REMEMBER

Good asthma control means having **NO** symptoms at all

If you have any symptoms you should speak to your doctor or asthma specialist as soon as possible



# Child Asthma Plan

Ages 4 - 11

Name:

Date:

Produced by  
London Babies, Children and Young People's Team

Every day:  
I am symptom free



My asthma is not  
controlled if...



I'm having an asthma attack  
and need to see a  
doctor now if...



### Preventer Inhaler

I need to take my preventer inhaler every day

It is called:

\* needs a spacer

and its colour is:

My best peakflow measure is:

 l/min

I take  puff/s of my preventer inhaler in the morning and  puff/s at night. I do this every day even if my asthma's OK

Other asthma medicines I take every day:

### Rescue Inhaler

My rescue inhaler helps when I am wheezy or coughing, finding it harder to breathe, or my chest hurts. I should not need it regularly.

It is called:

\* needs a spacer

and its colour is:

I take  puff/s when needed

I wheeze, cough, my chest hurts, or it's hard to breathe **or**

I regularly need my rescue inhaler one or more times a week **or**

If my asthma is stopping me doing sport or other activity **or**

I'm waking up at night because of my asthma (this is an important sign and I will book a next day appointment with my GP or nurse) **or**

My peakflow measure falls below 80%:

 l/min

### So I need to...

Take 2 puffs of my rescue inhaler, one puff at a time.

After 5-10 minutes, if I still have symptoms repeat this until I have had up to 6 puffs.

### I should feel much better

This should last at least 4 hours.

I will call my GP to arrange an appointment today or tomorrow

If I don't feel better, or my  
symptoms return within 4 hours,  
move to the next section

My symptoms aren't **COMPLETELY** better after 6 puffs of my rescue inhaler **or**

I need my rescue inhaler again in less than four hours **or**

My peakflow measure falls below 60%:

 l/min

I also need to take up to 10 puffs of  
my rescue inhaler, one puff at a  
time.

If my symptoms aren't completely better after 10 puffs

I will call 999 and tell them I'm  
having an asthma attack and it's  
not controlled by 10 puffs of my  
rescue inhaler

I also need to...



Sit up - don't lie down. Try to keep calm.

Take one puff of my rescue inhaler. Then repeat every 60 seconds.

If the ambulance has not arrived after 10 minutes, **contact 999 again immediately.**